THE OHIO STATE UNIVERSITY WEXNER MEDICAL CENTER

Medical News Public Affairs and Media Relations 614-293-3737 The James

THE OHIO STATE UNIVERSITY COMPREHENSIVE CANCER CENTER



Media Contact: Amanda J. Harper OSUCCC – James Media Relations Office: 614-685-5420 Cell: 513-659-6847 Amanda.Harper2@osumc.edu

NOTE TO EDITORS: High quality b-roll and images associated with this story are available for download at <u>http://bit.ly/2oP4DAt</u> (password: reconstruction).

EMBARGOED FOR RELEASE: May 3, 2017, 11:01 AM ET

Half of Breast Cancer Patients Pursue Reconstructive Surgery Without Understanding of Risks

New data show most mastectomy patients choose treatment that does not align with personal goals

COLUMBUS, Ohio – More than half of breast cancer patients (57 percent) undergoing mastectomy lack the necessary medical knowledge to make a high-quality decision about reconstructive surgery that aligns with their personal goals, suggesting a trend toward overtreatment, according to a new study conducted at <u>The Ohio State University Comprehensive Cancer Center – Arthur G. James</u> <u>Cancer Hospital and Richard J. Solove Research Institute</u> (OSUCCC – James).

"High-quality" decisions were defined as those that demonstrated adequate medical knowledge of treatment choices -- including associated risks – and that also matched with the patient's specific goals and preferences for choosing whether or not to pursue reconstructive surgery.

Researchers say shared decision-making tools are needed to help women make decisions based on a full understanding of treatment choices and associated risks alongside their personal goals for surgery.

Researchers report the findings online first in the medical journal JAMA Surgery May 3, 2017.

Study Design and Methods

In this observational, single-institution study, researchers sought to evaluate the quality of 126 adult breast cancer patients' decisions about breast reconstruction after mastectomy. All patients had

stage I-III invasive ductal/lobular breast cancer, ductal carcinoma in situ (DCIS) or were having preventive mastectomies, and the majority (73 percent) had early-stage disease.

Researchers measured study participants' medical knowledge about mastectomy and mastectomy with reconstruction -- for example, effects of surgery on appearance and associated risks. They also measured individual preferences of what mattered most to patients. Key preference factors included breast appearance/shape post treatment, length of recovery time and risk for complications.

"We found that less than half of the women had adequate medical knowledge about breast reconstruction and made a choice that aligned with their personal preferences. This is very concerning to us, because it means that some women did not get the treatment they truly preferred, and quite a few had more treatment than they preferred," says <u>Clara Lee</u>, MD, principal investigator of the study and a breast reconstructive surgeon at The OSUCCC – James. Lee holds a dual associate professor appointment in the colleges of medicine and public health at Ohio State.

"Many women were quite concerned about complication risks, but they didn't actually know how high the risk was. This may explain some of the overtreatment that we saw," she adds.

Researchers found that only 43 percent of the patients in the study demonstrated an understanding of at least half of the important facts about reconstruction and made a choice that was consistent with their preferences. Understanding of surgical complications was particularly low, with only 14 percent of patients demonstrating strong knowledge of associated risks.

"As breast cancer providers, we need to talk about the pros and cons of surgery to help women make treatment choices. Shared decision-making between the surgeon and patient would be particularly useful for this decision. We need to connect patients with decision aids to help them really think through what is most important to them," Lee adds.

Collaborators in this National Cancer Institute-funded study include Allison Deal, MD, and Ruth Huh, BA, of Lineberger Comprehensive Cancer Center at University of North Carolina Chapel Hill; Michael Pignone, MD, MPH, of University of Texas at Austin; and Peter Ubel, MD, of Duke University.

Patient-Doctor Communication Research Underway

Lee and colleagues in Ohio State's colleges of engineering, communication and public health are working on a study to evaluate treatment decisions in early-stage breast cancer patients to assess how communication with their providers affects their decision-making. This ongoing study examines patients' knowledge, preferences, and expectations about future well-being. Information from this study is expected to help clinicians develop tools to aid patients in making an informed decision about their care.

-30-

About the OSUCCC – James

The Ohio State University Comprehensive Cancer Center – Arthur G. James Cancer Hospital and Richard J. Solove Research Institute strives to create a cancer-free world by integrating scientific research with excellence in education and patient-centered care, a strategy that leads to better methods of prevention, detection and treatment. Ohio State is one of only 47 National Cancer Institute (NCI)-designated Comprehensive Cancer Centers and one of only a few centers funded by the NCI to conduct both phase I and phase II clinical trials on novel anticancer drugs sponsored by the NCI. As the cancer program's 308-bed adult patient-care component, The James is one of the top cancer hospitals in the nation as ranked by *U.S. News & World Report* and has achieved Magnet designation, the highest honor an organization can receive for quality patient care and professional nursing practice. At 21 floors and with more than 1.1 million square feet, The James is a transformational facility that fosters collaboration and integration of cancer research and clinical cancer care. Learn more at <u>cancer.osu.edu</u>.